

## Is there a medicine to treat the flu?

Antiviral drugs can treat flu illness. They can make people feel better and get better sooner and may prevent serious flu complications, like pneumonia, for example, that can lead to hospitalization and even death. These drugs are different from antibiotics, but they also need to be prescribed by a doctor. They work best when started during the first 2 days of illness. It's very important that antiviral drugs be used early to treat the flu in people who are very sick (for example people who are in the hospital) or people who are at greater risk of having serious flu complications. Other people with flu illness may also benefit from taking antiviral drugs. These drugs can be given to children and pregnant women.

## What are some of the other ways I can protect my child against the flu?

In addition to getting vaccinated, take – and encourage your child to take – everyday steps that can help prevent the spread of germs.

This includes:

- ▶ Cover coughs and sneezes with a tissue. Throw the tissue in the trash after you use it.
- ▶ Stay away from people who are sick.
- ▶ Wash hands often with soap and water. If soap and water are not available, use an alcohol-based hand rub.
- ▶ Avoid touching your eyes, nose and mouth. Germs spread this way.
- ▶ If someone in the household is sick, try to keep the sick person in a separate room from others in the household, if possible.
- ▶ Keep surfaces like bedside tables, surfaces in the bathroom, kitchen counters and toys for children clean by wiping them down with a household disinfectant according to directions on the product label.
- ▶ Throw tissues and other disposable items used by sick persons in your household in the trash.

These everyday steps are a good way to reduce your chances of getting all sorts of illnesses, but a yearly flu vaccine is always the best way to specifically prevent the flu.



## What should I use for hand washing?

Washing hands with soap and water (for as long as it takes to sing the "Happy Birthday" song twice) will help protect against many germs. If soap and water are not available, use an alcohol-based hand rub.

## IF YOUR CHILD IS SICK

### What can I do if my child gets sick?

Talk to your doctor early if you are worried about your child's illness.

If your child is 5 years and older and does not have other health problems and gets flu-like symptoms, including a fever and/or cough, consult your doctor as needed and make sure your child gets plenty of rest and drinks enough fluids.

If your child is younger than 5 years (and especially younger than 2 years) or of any age with a long term health condition (like asthma, a neurological condition, or diabetes, for example) and develops flu-like symptoms, they are at risk for serious complications from the flu. Ask a doctor if your child should be examined.

### What if my child seems very sick?

Even children who have always been healthy before or had the flu before can get very sick from the flu.

Call for emergency care or take your child to a doctor right away if your child of any age has any of the warning or emergency signs below:

- ▶ Fast breathing or trouble breathing
- ▶ Bluish or gray skin color
- ▶ Not drinking enough fluids (not going to the bathroom or making as much urine as they normally do)
- ▶ Severe or persistent vomiting
- ▶ Not waking up or not interacting
- ▶ Being so irritable that the child does not want to be held
- ▶ Flu-like symptoms improve but then return with fever and worse cough
- ▶ Has other conditions (like heart or lung disease, diabetes, or asthma) and develops flu symptoms, including a fever and/or cough.

### Can my child go to school, day care or camp if he or she is sick?

No. Your child should stay home to rest and to avoid giving the flu to other children or caregivers.

### When can my child go back to school after having the flu?

Keep your child home from school, day care or camp for at least 24 hours after their fever is gone. (Fever should be gone without the use of a fever-reducing medicine.) A fever is defined as 100°F (37.8°C) or higher.

For more information, visit [www.cdc.gov/flu](http://www.cdc.gov/flu) or [www.flu.gov](http://www.flu.gov) or call 800-CDC-INFO